

used. The mobile phase consisted of water (solvent A) and methanol (solvent B), both containing 2 mM ammonium acetate. The analytes were eluted with a linear gradient of 30-90 % solvent B from 0 to 6 min, with a constant flow rate of 0.300 mL/min. TNT, and the TNT metabolites 2-ADNT and 4-ADNT were analyzed by multiple reaction monitoring (MRM) of the compounds respective parent and daughter molecules formed by electrospray ionization (ESI) in negative mode, in principle as described by Lotufo et al. (2016). Quantification was performed with 2, 4, 6-TNT ($^{13}\text{C}_7$) as internal standard. TNT was monitored at m/z 226 $[\text{TNT-H}]^-$ (parent ion) and m/z 197 $[\text{TNT-H-NO}]^-$ (daughter ion); the latter was used for quantification. 2-ADNT and 4-ADNT coeluted on the separation column, and had identical parent ions $[\text{M-H}]^-$ of m/z 196. However, the intensity of the daughter ions differed which enabled us to differentiate between the two compounds. The daughter ions selected for quantitation of 2-ADNT and 4-ADNT was m/z 136 $[\text{M-H-2NO}]^-$ and m/z 149 $[\text{M-H-HNO}_2]^-$, respectively. The fragmentation patterns of 2-ADNT and 4-ADNT revealed some presence of m/z 149 (~30%) and m/z 136 (~15%), respectively, implying that the quantified results are somewhat overestimated. The results of 2-ADNT and 4-ADNT must therefore be considered semi quantitative.

2.8. Data analysis.

Bioconcentration factors (BCF) in the depuration experiment were estimated as $\mu\text{g } ^{14}\text{C-TNT}$ equivalents per kg tissue divided by $\mu\text{g } ^{14}\text{C-TNT}$ equivalents per L of water at the same time point as the fish were sampled. BCF in the second experiment were estimated as $\mu\text{g TNT}$ per kg tissue divided by $\mu\text{g TNT}$ per L in the exposure waters at the end of the experiment. Elimination half-time was estimated from non-linear regression one-phase exponential decay. Descriptive statistics, non-linear

regression analysis (one phase exponential decay), ANOVA and mathematical calculations were computed in GraphPad Prism 5 or Excel 2007.

3. Results

3.1. TNT in the exposure water

The radioactivity in the water decreased during the first 10 h followed by a subsequent increase (Fig 1). The radioactivity in the water in the depuration tank increased during the first 20 h after the transfer of the fish (Fig 1).

In the second experiment, water was analyzed for TNT, 2-ADNT and 4-ADNT. Water from the respective exposure tanks were analyzed at the start of the experiment and after approximately 8 h, 24 h and 48 h showing a gradually a decrease of TNT in the water as a function of time. After 48 h the reduction in the TNT concentration ranged from 78% in the 100 µg/L tank to 16% in the water from the Lake Maridalsvannet (Table 1). The concentrations of 2-ADNT and 4-ADNT increased during the first 24 h followed by a decrease (Table 2), except in the water from the Lake Maridalen.

3.2. Uptake and excretion of TNT in fish

In the gills, blood, liver, kidney, muscle and the brain it was a rapid increase in the uptake of ^{14}C -TNT with a maximum tissue concentration after 6 hours (12 hours in the brains) (Fig 2). After 6 hours (12 hours in the brains) the concentration of ^{14}C -TNT in the organs started to decrease, even though the fish was still exposed to ^{14}C -TNT in the water. The radioactivity in the gallbladder increased during the whole experiment, reaching a maximum after 55 hours, 7 hours after the transfer to the clean water (Fig 3). A similar trend was observed in the intestines. The excretion of ^{14}C -TNT in the fish started before they were transferred to the clean water. Estimated elimination half-time of ^{14}C -TNT in the fish from the time the depuration started ranged from 8 hours in the liver to 22 hours in the kidney (Fig 1S, Supplementary materials). It was not feasible

to make a good estimation of the elimination half-life of ^{14}C -TNT in the fish after the transfer to the clean water since the concentration in the different organs already was near the measured minimum levels.

BCF were estimated after 6 hours of exposure when maximum tissue concentrations were observed and after 48 hours when the fish were transferred to the clean water. With the exception of the gallbladder the BFCs were highest in the gills, kidney, livers and intestine representing the first target of exposure and organs of excretion and metabolism (Table 3). The BCFs in these organs decreased considerable after 6 hours of exposure. The BCF of ^{14}C -TNT in the gall bladder increased from 404 after 6 hours of exposure to nearly 2000 after 48 hours of exposure.

In fish exposed to unlabeled TNT, TNT, 2-ADNT and 4-ADNT were found in the muscle tissue. Only 2-ADNT and 4-ADNT were found in the bile samples (Table 4 and 5). Bioconcentration factors were estimated based on the concentration in the tissue and the concentrations in the water and was between 4 and 6 (Table 4)

3.3 Effects of increasing TNT concentrations on Atlantic salmon

One fish in each of the two groups exposed to the highest concentration (1 mg/L) died during the experiment. During the dissection procedure it was observed that all the fish exposed to the highest concentrations had severed hemorrhages in the dorsal muscle tissue near the spine (Fig 2S). All the fish exposed to the highest concentrations would probably have died if the exposure time had been extended. There were no visible signs of any injuries or behavioral abnormalities on the fish exposed to the lower concentrations. With the exception of the groups exposed to the highest concentrations there were no significant effects on any of the blood physiology parameters that were analyzed. Severe effects on blood physiology was observed in

the two groups exposed to 1 mg/L of TNT (Table 6 and 7). In fish exposed to synthetic lake water, increased levels of glucose, urea and HCO_3 were observed, while hematocrit and the levels of Cl and hemoglobin (Table 6) decreased. In the fish exposed to TNT in the water from Lake Maridalsvannet it was observed an increase in urea and glucose, and a decrease in hematocrit and the levels of hemoglobin (Table 7).

4. Discussion

4.1. TNT in the exposure water

The experiment with use of ^{14}C -labeled TNT showed a loss of approximately 25% of the added ^{14}C -TNT from the water during the first 12 hours followed by a subsequent increase in the radioactivity (Fig 1). The subsequent increase in the radioactivity is probably attributed to excretion from the fish, bearing in mind that the depuration started before the fish were transferred to the clean water. At the end of the experiment a reduction in total radioactivity of 12% was observed. The loss may primarily be attributed to uptake and sorption in the fish, but may also be due to adsorption to the exposure tank walls made of polyethylene which has hydrophobic properties. In the second experiment with unlabeled TNT the findings were more complex. In all the exposure groups a reduction in the TNT concentrations were observed in the water, ranging from 60 to 80% reductions in the groups exposed to 1-100 $\mu\text{g/L}$ and a 15-40% reduction in the groups exposed to 1 mg/L (Table 1). The loss of TNT during the experiment indicated uptake and/or metabolism in the fish, attachment to the tank walls, and/or decomposition in the water. Apparently the reduction in the TNT concentration was less pronounced in the water from the Lake Maridalsvannet, of which it was observed a loss of approximately 15%. Water from Lake Maridalsvannet was also used in the first experiment, which had a similar loss of radioactivity during the experiment. Unlike the synthetic water, the water from Lake Maridalsvannet contain organic materials (approximately 4.5 mg/L), which TNT and metabolites may be associated with. It has previously been shown that TNT may be adsorbed to organic materials in soil, such as humic acids (Esteve-Nunez et al., 2001; Eriksson et al., 2004). The metabolites of TNT, 2-ADNT and 4-ADNT, were detected in the water. The amount of metabolites, could however, not account for the loss of

TNT, indicating formation of considerable amount of unknown decomposition products, or adsorption to the tank walls and to the fish. In the ^{14}C -TNT experiment a considerable fraction of radioactivity retained in the different organs at the end of the study (Fig 2). This observation indicates that some of the losses of TNT in the water may be due to non-extractable TNT-compounds adsorbed to fish tissues. Previous studies on effects of TNT on aquatic organisms have shown similar losses of TNT in the exposure water (Conder et al., 2004; Lotufo et al., 2010; Yoo et al., 2006). Possible decomposition products other than the ADNT, may be diamino nitrotoluenes, tetranitroazoxytoluenes, nitrobenzoic acids and nitrogen free TNT transformation products (Conder et al., 2004; Preiss et al., 2005; Esteve-Nunez et al., 2001; Carpenter et al., 1978; Kaplan and Kaplan, 1982; Smith et al., 2015).

4.2. Uptake and excretion of ^{14}C -TNT in fish

A rapid uptake of ^{14}C -TNT was observed in the gills, blood, liver, kidney, muscle and the brain after 6 hours of exposure. In the brains maximum concentration was reached after 12 hours of exposure. The delayed uptake in brain demonstrate that the ^{14}C -TNT compounds passes the blood brain barrier. The maximum concentration of ^{14}C -TNT in the fish brain was similar to the concentrations detected in other tissues. After 6 hours in six of the tissues, and 12 hours in the brains, the concentration of ^{14}C -TNT started to decrease, even though the fish was still exposed in the water. Since the Atlantic salmon was not fed, and do not drink water, the site of TNT uptake must be the gills. The mechanism behind the reduced uptake after the initial rapid uptake is unclear, but it might be linked to negative effects on important processes at the gill surface and ion exchange mechanisms, demonstrated by the negative effects on osmoregulation and general physiology. Changes in mucus secretion or mucus quality

might also be an explanation, although not measured in our experiment. Sensini et al. (2008) exposed European eel to TNT in the water for concentrations ranging from 0.5 – 2.5 mg/L. In addition to lesions on the gills, such as oedema and vascular congestion, they observed mucus hypersecretion, which may be a response to an exposure to a xenobiotic leading to reduced uptake.

BCF were estimated after 6 hours of exposure to ^{14}C -TNT when maximum uptake was observed and after 48 hours when the exposure period was terminated. With the exception of the gall bladder the BCFs were highest in the gills, kidney, livers and intestine, representing first target of exposure and organs of excretion and metabolism (Table 3). The BCFs decreased considerable after 48 hours showing that ^{14}C -TNT is not particular prone to bioaccumulation and that it is readily excreted. The BCF of ^{14}C -TNT in the gall bladder increased from 404 L/kg after 6 hours of exposure to nearly 2000 L/kg after 48 hours of exposure. Apparently the ^{14}C -TNT was concentrated into the gall bladder showing that the fish primarily excrete TNT and potential metabolites through the bile. In most organs, the concentrations of ^{14}C -TNT were close to the measured minimum when the fish were transferred to clean water and elimination half-life could not be calculated. Previous studies have estimated elimination half-life in different fish species of an hour or less (Ownby et al., 2005, Lotufo and Lydy , 2005; Yoo et al., 2006). The elimination half-time of TNT in the fish was, however, estimated from the time the depuration started and ranged from 8 hours in the liver and 22 hours in the kidney (Fig 1S).

At the termination of the experiment there were still some residual ^{14}C -TNT left in the fish organs (Fig 2) indicating a slow elimination of a portion of the accumulated ^{14}C -TNT. This portion may be the so-called unextractable fraction, which may be biotransformation products covalently bound to tissue proteins and other

macromolecules as discussed by Ownby et al, (2005). The reductive metabolism of TNT to aminonitrotoluenes involves formation of several reactive metabolites, which have shown to form adducts with thiol groups (Leung et al., 1995).

4.3. Accumulation of TNT, 2-ADNT and 4-ADNT in the fish

TNT and its metabolites 2-ADNT and 4-ADNT, were detected in the muscle tissue (Table 4 and 5). The concentrations of TNT in the muscle tissue increased with the exposure concentrations. The calculated BCFs of TNT in muscle tissue were in the range of 4-6 L/Kg (Table 4) confirming that TNT is not particularly prone to bioaccumulation and that it is readily excreted. With the ¹⁴C-TNT study in mind, however, it is reasons to believe that the accumulated concentrations of TNT in the effect study were higher earlier in the experiment, maybe by a factor of 3-5. The concentrations of the ADNT-metabolites were similar as the concentration of the parent molecule, indicating that a considerable fraction of the parent molecule is metabolized.

The ¹⁴C-TNT experiment showed that the radio labeled TNT accumulated in the gall bladder. This was confirmed by chemical analysis of the bile of the TNT exposed fish. We did, however, only detect the presence of 2-ADNT and 4-ADNT and not the parent molecule (Table 5). The bile in our study was treated with β -glucuronidase during preparation. Previously Eek et al (2005) could not recover any TNT and TNT metabolites in the bile from fish orally exposed to TNT (100-400 mg/kg body weight). By treating the bile with β -glucuronidase in order to hydrolyze glucuronide conjugated TNT and TNT metabolites it was shown that considerable portion of the TNT was excreted as glucuronides. Eek et al., (2005) found that that the concentrations of the metabolites in the bile were a factor of approximately 10 higher

than the parent compound, which may explain the reason why we did not detect TNT in the bile in our study. In mammals, comparative studies have shown that the major excretion pathway of TNT is through urine primarily as the glucuronide conjugates (El-Hawari et al., 1981; Talmage et al., 1999). In fish it appears that a major portion of the TNT is metabolized followed by a phase II conjugation with glucuronides and biliary excretion.

In order to identify potential exposure of TNT near dumping sites of ammunition it may be reasonable to analyze the fish bile. In this study TNT metabolites were detected in bile from fish exposed to 10 µg/L. With improved analytical techniques it will probably be possible to detect TNT and breakdown products in bile from fish exposed to lower concentrations (e.g. Ochsenberg et al., 2008). Due to dilution of munition residues in the seawater there are often no detection or only trace amounts of energetics residues that are detected near ammunition dump sites (e.g. Darrach et al., 1998; Ampleman et al., 2004; Ochsenberg et al., 2008; Rosland et al., 2010; University of Hawaii, 2011). In close vicinity of the munitions, however, considerable concentrations have been reported. At the Bedford Basin, Halifax Nova Scotia, large amounts of ammunition from WWI and WWII are dumped (Rodacy et al., 2001). Water was sampled in close vicinity of the ammunition (≤ 1 m), and at one site it was found 14 µg/L TNT, 123 µg/L 4-ADNT and 108 µg/L 2-ADNT in the water. Barton and Porter (2004) performed a survey at the eastern end of Isla de Vieques, Puerto Rico which were used as a naval gunnery and bombing range between 1943 and 2003. In seawater inside a bomb it was found 66.4-105 mg/L TNT showing considerable dissolution of the energetic materials. Approximately one meter from the bomb the concentrations had dropped to 17.7 and 7.9 µg/L. In order to assess risk of exposure from leakage of very low concentrations of munitions residues from

dumping sites or training areas, other sampling strategies, such as passive samplers, may be more convenient (Rosen et al., 2016; Belden et al., 2015) or biomonitoring with transplanted blue mussels (Strehse et al., 2017).

4.4. Effects of TNT on fish

In the two groups exposed to the highest concentration mortality was observed, and severe hemorrhages in the muscle tissue at the dorsal fin near the spine (Fig 2S). The effect of TNT on fish is species dependent with reported acute lethal concentrations ranging from 1-5 mg/L, of which species like fathead minnow and channel catfish appear less sensitive than rainbow trout (Liu et al., 1983). A previous study reported a mean LC₅₀-concentration of 1.15 mg/L on rainbow trout exposed for 96 h under static condition (Liu et al., 1983). Compared with other aquatic organisms fish appears to be one of the most sensitive taxa to TNT intoxication (Liu et al., 1983; Ribeiro et al., 2012; Lotufo et al., 2013) and Atlantic salmon appears to be a particular sensitive species. Leffler et al., (2014) exposed alevins of Atlantic salmon to TNT wastewater for 40 days. In the high exposure group (2.1 mg/L TNT) they observed approximately 25% mortality after 14 days and 100% mortality after 40 days. In the group exposed to 0.41 mg/L they observed approximately 30% mortality after 40 days. In comparison to our experiment it is apparent that alevins are somewhat less susceptible than the juveniles.

The effect of TNT on mammals has been thoroughly studied and it is reasons to believe that similar mechanisms of toxicity are involved in the fish. One of the most pronounced symptoms of severe TNT intoxication in mammals is anemia (Crawford 1954; Dilley et al., 1982; Ryon and Ross, 1990; Bradley 2011). The symptoms are caused either by hemolysis of the red blood cells (as we saw in our experiment), by a

destruction of the hemoglobin in the cells, or by a chemical change in the hemoglobin of the red blood corpuscles, forming mixtures of methhemoglobin, NO-hemoglobin, and sulphhemoglobin with a consequent damage of the oxygen-carrying capacity. The analysis of the physiological parameters in blood confirmed severe effects in our experiment (Table 6 and 7). These effects on blood were only observed in the fish exposed to the highest concentration of TNT. Both the effect on morphology and on blood physiology appeared independent of the water quality. The fish exposed to TNT in the synthetic water had, however, effects on two more blood parameters, but it is reason to believe that similar effect eventually had happened also in the fish exposed to the TNT in the water from Lake Maridalsvannet. Eek et al., (2003) who exposed rainbow trout perorally to TNT (100 mg/kg - 400 mg/kg) for 48 h did not report any morphological changes but observed increased levels of methhaemoglobin and increased numbers of lymphocytes and leukocytes, which is in accordance with previous studies on mammals and the present study. The effect observed on the fish is probably due to formation of reactive metabolites of TNT. In mammals the nitro groups on the TNT undergo a reductive metabolism into aminonitrotoluenes. The reduction of TNT is a stepwise reaction with the conversion of TNT to the intermediates nitrosodinitrotoluene and hydroxylaminodinitrotoluene. The conversion of hydroxylaminodinitrotoluene to aminodinitrotoluene is a rate limiting step which may cause an accumulation of the substance. Nitrosodinitrotoluene and hydroxylaminodinitrotoluene are reactive metabolites which may covalently bind to macromolecules in the cells (Levine et al., 1990; Michels et al., 1994; Leung et al., 1995; Bakhtiar and Leung, 1997; Esteve-Nunez, et al., 2001). Apparently, in the high exposure group in our experiment a level of metabolites was reached that was detrimental for the fish.

4.5. Conclusions

The present study have shown that TNT is taken up primarily over the gills and rapidly excreted from fish via the bile. After 6 hours (12 hours in the brains) the concentration of TNT in the organs started to decrease, even though the fish was still exposed to TNT in the water. The depuration experiment showed that the BCF for TNT-equivalents were substantially higher in the gall bladder, likely due to presence in the bile, compared to the other organs. In order to identify potential exposure of TNT near dumping sites of ammunition it may be reasonable to analyze the fish bile. Mortality was observed amongst fish exposed to 1 mg/L TNT. Considerable amount of the TNT metabolites 2-ADNT and 4-ADNT, were detected in the muscle tissue of the fish, and severe hemorrhages in the dorsal muscle tissue near the spine was observed. Effects on blood parameters such as of glucose, urea, hematocrit and hemoglobin were observed in the fish exposed to the highest concentration (1 mg/L). Large amount of ammunition have been dumped in the oceans of the world during the last century, but due to dilution of munitions residues in the seawater there is less likely that fish are particular vulnerable to exposure. Benthic organisms are probably at highest risk of exposure (Pascoe et al., 2010; Voie and Mariussen, 2017). However, in an area with little exchange of water, like a small lake or a threshold fjord, or in close vicinity of the ammunition, dissolved TNT and other ammunition residues may reach toxic levels.

Acknowledgements

This study was funded by the Ministry of Defence (FFI Project No. 119701 and 130001). Some funding were granted from the DAIMON project (financed under the Interreg Baltic Sea Programme 2014-2020) to finish the manuscript. The authors want to thank Dr. Merethe Kleiven, Ms. Marit Nandrup Pettersen from NMBU, and Ms Ida Vaa Johnsen from FFI for helping with the dissection of the TNT exposed fish, and Ms Anita Larsen and Ms Anne Myhran from FFI for helping with the extraction of water and muscle tissue for the chemical analysis.

Conflict of interest statement

The authors declare that there are no conflicts of interest.

References

- Ampleman G., Faucher, D., Thiboutot, S., Hawari, J., Monteil-Rivera F., (2004).
Evaluation of underwater contamination by explosives and metals at Point Amour
Labrador and in the Halifax Harbour area. Technical report DRDC Valcartier TR
2004-124. Valcartier, Quebec, Canada: Defense Research and Development
Canada.
- Bakhtiar, R., Leung, K.H., 1997. Covalent binding of 2,4,6-trinitrotoluene to human
hemoglobin. Evidence for protein adducts probed by electrospray ionization mass
spectrometry. *Rapid Commun. Mass Spectrom.* 11, 1935-7.
- Barton, J.V., Porter, J.W (2004). Radiological, Chemical, and Environmental Health
Assessment of the Marine Resources of the Isla de Vieques Bombing Range, Bahia
Salina del Sur, Puerto Rico. CEO, Underwater Ordnance Recovery, Inc and the
University of Georgia, Athens.
- Belden, J.B., Lotufo, G.R., Biedenbach, J.M., Sieve, K.K., Rosen, G., 2015.
Application of POCIS for exposure assessment of munitions constituents during
constant and fluctuating exposure. *Environ. Toxicol. Chem.* 34, 959-967.
- Bradley, M.D., 2011. 2,4,6-Trinitrotoluene (TNT) air concentrations, hemoglobin
changes, and anemia cases in respirator protected TNT munitions demilitarization
workers. *Int. Arch. Occup. Environ. Health.*, 84, 239-250.
- Bradley, P.M., Chapelle, F.H., Landmeyer, J.E., Schumacher J.G., 1994. Microbial
transformation of nitroaromatics in surface soils and aquifer materials. *Appl.*
Environ. Microbiol. 60, 2170-2175.
- Carpenter, D.F., McCormick, N.G., Cornell, J.H., Kaplan, A.M. 1978. Microbial
transformation of ¹⁴C-labelled TNT in an activated-sludge system. *Appl.*
Environment. Microbiol. 35, 949-954.

- Conder, J.M., La Point, T.W., Bowen, A.T., 2004. Preliminary kinetics and metabolism of 2,4,6-trinitrotoluene and its reduced metabolites in an aquatic ologochaete. *Aq. Tox.* 69, 199-213.
- Crawford, M.A. 1954. Aplastic anaemia due to trinitrotoluene intoxication. *Br Med. J.* 2, 430-437.
- Darrach, M.A., Chutjian, A., Plett, G.A., 1998. Trace Explosives Signatures from World War II Unexploded Undersea Ordnance. *Environ. Sci. Technol.* 32, 1354–1358.
- Dilley, J.V., Tyson, C.A., Spangord, R.J., Sasmore, D.P., Newell, G.W., Dacre, J.C., 1982. Short-term oral toxicity of 2,4,6-trinitrotoluene in mice, rats, and dogs. *J Toxicol Environ. Health.*, 9, 565-85.
- Ek, H., Dave, G., Birgersson, G., Förlin, L. 2003. Acute effects of 2,4,6-trinitrotoluene on heamatology parameters and hepatic EROD-activity in rainbow trout (*Oncorhynchus mykiss*). *Aquat. Ecosyst. Health Manag.* 6, 415-421.
- Ek, H., Dave, G., Sturve, J., Almroth, B.C., Stephensen, E., Förlin, L., Birgersson, G. 2005. Tentative biomarkers for 2,4,6-trinitrotoluene (TNT) in fish (*Oncorhynchus mykiss*). *Aquat Toxicol.* 72, 221-230.
- El-hawari, A.M., Hodgson, J.R., Wnston, J.M., Sawyer, J.M., Hainje, M., Lee, C.C. 1981. Species Differences in the Disposition and Metabolism of 2,4,6-Trinitrotoluene as a Function of Route of Administration Final report. AD A114025. US army Mediac Research and Development Command, Fort Detrick, MD.
- Eriksson, J., Frankki, S., Shchukarev, A., Skyllberg, U., 2004. Binding of 2,4,6-trinitrotoluene, aniline, and nitrobenzene to dissolved and particulate soil organic matter. *Environ. Sci. Technol.* 38, 3074-3080.

- Esteve-Núñez, A.1., Caballero, A., Ramos, J.L., 2001. Biological degradation of 2,4,6-trinitrotoluene. *Microbiol. Mol. Biol. Rev.* 65, 335-352.
- ICP Waters 2010. Chapter 11. Trace metals and persistent organic pollutants in fish. Page 69-81. In: ICP Waters Programme Manual 2010. NIVA Report 6074-2010, ICP Waters Report 105/2010 (<http://www.icp-waters.no/publications/#icpwmanual>)
- Kaplan, D.L., Kaplan, A.M., 1982. Termophilic biotransformation of 2,4,6-trinitrotoluene under simulated composting conditions. *Advan. Appl. Biotechnol.* 44, 757-760.
- Leffler, P., Brännäs, E., Ragnvaldsson, D., Wingfors, H., Berglind, R., 2014. Toxicity and accumulation of trinitrotoluene (TNT) and its metabolites in Atlantic salmon alevins exposed to an industrially polluted water. *J. Toxicol. Environ. Health A.* 77, 1183-1191.
- Leung, K.H., Yao, M., Stearns, R., Lee Chiu, S.H., 1995. Mechanism of bioactivation and covalent binding of 2,4,6-trinitrotoluene. *Chemico. Biol. Inter.* 97, 37-51.
- Levine, B.S., Furedi, E.M., Gordon, D.E., Barkley, J.J., Lish, P.M., 1990. Toxic interactions of the munitions compounds TNT and RDX in F344 rats. *Fundament. Appl. Toxicol.* 15, 373-380.
- Levsen, K., Mussmann, P., Berger-Preiss, E., Preiss, A., Volmer, D., Wunsch, G., 1993. Analysis of nitroaromatics and nitramines in ammunition waste water and in aqueous samples from former ammunition plants and other military sites. *Acta Hydrochim. Hydrobiol.*, 21 153-166.
- Liu, D.H.W., Spangord, R.J., Bailey, H.C., Javitz, H.S., Jones D.C.L., 1983. Toxicity of TNT wastewaters to aquatic organisms. Final report, Volume 1. Acute toxicity of LAP wastewater and 2,4,6-trinitrotoluene. Report no AD-A142 144, US Army medical research and development command Fort Detrick, Frederick, MD 21701.

- Lotufo, G.R., Belden, J.B., Fisher, J.C., Chen, S.F., Mowery, R.A., Chambliss, C.K., Rosen, G., 2016. Accumulation and depuration of trinitrotoluene and related extractable and nonextractable (bound) residues in marine fish and mussels. *Environ Pollut.* 210, 129-136.
- Lotufo, G.R., Biedenbach, J.M., Sims, J.G., Chappell, P., Stanley, J.K., Gust, K.A. (2015). Bioaccumulation kinetics of the conventional energetics TNT and RDX relative to insensitive munitions constituents DNAN and NTO in *Rana pipiens* tadpoles. *Environ. Toxicol. Chem.* 34, 880-886.
- Lotufo G. R., Rosen, G. Wild, W. Carton G., 2013. Summary of Review of the Aquatic Toxicology of Munitions Constituents. Technical report US Army Corps of Engineers, ERDC/EL TR-13-8, 2013.
- Lotufo, G.R., Blackburn, W.M., Gibson, A.B., 2010. Toxicity of trinitrotoluene to sheepshead minnows in water exposures. *Ecotoxicol. Environ. Saf.* 73, 718-726
- Lotufo, G.R., Lydy, M.J., 2005. Comparative toxicokinetics of explosive compounds in sheepshead minnows. *Arch. Environ. Contam. Toxicol.* 49, 206-214.
- Mahrosh, U., Kleiven, M., Meland, S., Rosseland, B.O., Salbu, B., Teien, H.C., 2014. Toxicity of road deicing salt (NaCl) and copper (Cu) to fertilization and early developmental stages of Atlantic salmon (*Salmo salar*). *J. Haz. Mat.* 280, 331-339.
- Marencic H., Nehring, S., 2009. Military Activities. Thematic Report No. 3.5. In: Marencic, H. & Vlas, J. de (Eds), 2009. Quality Status Report 2009. Wadden Sea Ecosystem No. 25. Common Wadden Sea Secretariat, Trilateral Monitoring and Assessment Group, Wilhelmshaven, Germany, 2009.
- Michels, J., Gottschalk, G., 1994. Inhibition of the lignin peroxidase of *Phanerochaete chrysosporium* by hydroxyl-amino dinitrotoluene, an early intermediate in the degradation of TNT. *Appl. Environ. Microbiol.* 60, 187-194..

- Ochsenbein, U., Zeh, M., Berset, J.D., 2008. Comparing solid phase extraction and direct injection for the analysis of ultra-trace levels of relevant explosives in lake water and tributaries using liquid chromatography-electrospray tandem mass spectrometry. *Chemosphere* 72, 974-980.
- OECD 1992. Test guideline 203: Fish acute toxicity test.
- Ownby, D.R., Belden, J.B., Lotufo, G.R., Lydy, M.J., 2005. Accumulation of trinitrotoluene (TNT) in aquatic organisms: part 1--Bioconcentration and distribution in channel catfish (*Ictalurus punctatus*). *Chemosphere* 58, 1153-1159.
- Pascoe, G.A., Kroeger, K., Leisle, D., Feldpausch, R.J., 2010. Munitions constituents: Preliminary sediment screening criteria for the protection of marine benthic invertebrates. *Chemosphere*, 81, 807-816.
- Preiss, A., Elend, M., Gerling, S., Tränckner, S., 2005. Analysis of highly polar compounds in groundwater samples from ammunition waste sites. Part 1- Characterization of pollutant spectrum. *Magn. Reson. Chem.* 43, 736-746.
- Ribeiro, E.N., Da Silva, F.T., De Paiva, T.C., 2012. Ecotoxicological evaluation of wastewater from 2,4,6-TNT production. *J Environ. Sci. Health A Tox. Hazard Subst. Environ. Eng.* 47, 184-191.
- Rosen G, Wild B, George RD, Belden JB, Lotufo GR (2016). Optimization and field demonstration of a passive sampling technology for monitoring conventional munition constituents in aquatic environments. *Mar. Technol. Soc. J.* 50, 23-32.
- Rosseland, B.O., Massabuau, J-C., Grimalt, J., Hofer, R., Lackner, R., Raddum, G., Rognerud, S., Vives, I. 2001. Fish ecotoxicology, The EMERGE fish sampling manual for live fish. The EMERGE Project (European Mountain lake Ecosystems: Regionalisation, diagnostic and socio-economic valuation). (<http://www.mountain-lakes.org/emerge/methods/29.pdf>)

- Rosslund HK, Johansen A, Karsrud TE, Parmer MP, Larsen A, Myran A, Nordås SV (2010). Contamination From Dumped Conventional Munitions in the Aquatic Environment - Preliminary Investigation. [Norwegian] FFI/Report-2010/00239, 2010.
- Ryon, M.G., Ross, R.H., 1990. Water quality criteria for TNT. Regul. Pharmacol. 11, 104-113.
- Sensini, C., Della Torre, C., Corsi, I., Focardi, S., 2008. First observations of histopathological effects of 2,4,6-trinitrotoluene (TNT) in gills of European eel *Anguilla anguilla* (Linnaeus, 1758): histopathological effects of 2,4,6-trinitrotoluene in gills of European eel. Cell Biol. Toxicol. 24, 621-628.
- Sims, J.G., Steevens, J.A. (2008). The role of metabolism in the toxicity of 2,4,6-trinitrotoluene and its degradation products to the aquatic amphipod *Hyalella azteca*. Ecotoxicol. Environ. Saf. 70, 38-46.
- Smith, R.W., Vlahos, P., Böhlke, J.K., Ariyaratna, T., Ballentine, M., Cooper, C., Fallis, S., Groshens, T.J., Tobias, C., 2015. Tracing the Cycling and Fate of the Explosive 2,4,6-Trinitrotoluene in Coastal Marine Systems with a Stable Isotopic Tracer, (15)N-[TNT]. Environ. Sci. Technol. 49, 12223-12231.
- Smock, L.A., Stoneburner, D.L., Clark, J.R., 1976. The toxic effects of TNT and its primary degradation products on two species of alga and the fathead minnow. Wat. Res., 10, 537-543.
- Spalding, R.F., Fulton, J.W., 1988. Groundwater munition residues and nitrate near grand island, Nebraska, USA. J. Contam. Hydrol. 2, 139-153.
- Strehse, J.S., Appel, D., Geist, C., Martin, H.J., Maser, E. (2017). Biomonitoring of 2,4,6-trinitrotoluene and degradation products in the marine environment with transplanted blue mussels (*M. edulis*). Toxicol. doi.org/10.1016/j.tox.2017.09.004.

- Talmage, S.S., Opresko, D.M., Maxwell, C.S., Welsh, C.J.E., Cretella, F.M., Reno, P.H., Daniel, F.B., 1999. Nitroaromatic munition compounds: environmental effects and screening values. *Rev. Environ. Contam. Toxicol.* 161, 1-156.
- Yoo, L.J., Lotufo, G.R., Gibson, A.B., Steevens, J.A., Sims, J.G., 2006. Toxicity and bioaccumulation of 2,4,6-trinitrotoluene in fathead minnow (*Pimephales promelas*). *Environ. Toxicol. Chem.* 25, 3253-3260
- University of Hawaii, 2014. Final environmental study: Ordnance Reef (HI-06), Wai'anae, O'ahu, Hawai'i. Contract No. N00024-08-D-6323. Prepared for the U.S. Army Corps of Engineers.
- US EPA 1996. Ecological effects test guidelines. OPPTS 850. 1075. Fish acute toxicity test, freshwater and marine EPA 712-C-96-118, United States Environmental Protection Agency
- Voie Ø.A., Mariussen, E., 2017. Risk Assessment of Sea Dumped Conventional Ammunition. *Propell. Explos. Pyrot.* 42, 98-105.

Table 1

Concentrations ($\mu\text{g/L}$) of TNT in the water as a function of time in the effect study with unlabeled TNT. The concentrations are based on chemical analyses of one sample collected at each time point. The numbers in brackets show portion of TNT in percent relative to the start concentrations. M = water from Lake Maridalsvannet.

Exposure groups ($\mu\text{g/L}$)	Measured concentrations, $\mu\text{g/L}$ (% of start conc.)			
	Start	8h	24h	48h
1	1.0	1.1 (108)	0.4 (42)	0.3 (31)
10	12.6	11 (87)	7.14 (57)	4.8 (38)
100	108	101 (94)	58 (54)	24 (22)
1000	1014	990 (98)	780 (77)	567 (56)
1000 (M)	1026	939 (92)	927 (90)	864 (84)

Table 2

Concentrations ($\mu\text{g/L}$) of the TNT metabolites, 2-ADNT and 4-ADNT, in the exposure water as a function of time in the effect study with unlabeled TNT. The concentrations are based on chemical analyses of one sample collected at each time point. M = water from Lake Maridalsvannet.

Exposure groups ($\mu\text{g/L}$)	2-ADNT ($\mu\text{g/L}$)				4-ADNT ($\mu\text{g/L}$)			
	Start	8h	24h	48h	Start	8h	24h	48h
1	< 0.1	< 0.1	< 0.1	< 0.1	< 0.1	< 0.1	< 0.1	< 0.1
10	< 0.1	0.2	0.4	0.3	< 0.1	0.4	1.1	1.4
100	0.35	1.5	2.4	1.8	0.5	2.9	8	7.4
1000	2.3	5.9	10	1.8	1.5	7.4	26	16
1000 (M)	< 0.1	0.4	12	17	0.4	1.0	9.8	16

Table 3

Estimated bioconcentration factors (BCF) (mean BCF (L/kg) \pm SD, n = 6) of radiolabeled ^{14}C -TNT in the different organs after 6 h and 48 h of exposure.

Organ	BCF (6 hours)	BCF (48 hours)
Gills	46 \pm 5.0	9.6 \pm 0.5
Blood	10 \pm 1.4	2.5 \pm 0.6
Liver	43 \pm 9.2	18 \pm 3.0
Kidney	57 \pm 9.4	13 \pm 1.3
Muscle	14 \pm 0.7	2.4 \pm 0.4
Brain	15 \pm 1.0	6.7 \pm 0.5
Intestine	41 \pm 3.9	94 \pm 18
Gall bladder	404 \pm 75	1933 \pm 354

Table 4

The concentrations of TNT and TNT metabolites in muscle tissue, and estimated BCF of TNT. The results are shown as mean concentrations (mg/kg) \pm SD from 4 fish.

Exposure ($\mu\text{g/L}$)	TNT (mg/kg)	BCF TNT (L/kg)	2-ADNT (mg/kg)	4-ADNT (mg/kg)
1	< 0.05	n.e. ^a	< 0.05	< 0.05
10	< 0.05	n.e.	< 0.05	< 0.05
100	0.15 \pm 0.01	6.3 \pm 0.6	0.04 \pm 0.01	0.08 \pm 0.05
1000	2.9 \pm 0.6	5.2 \pm 1.0	0.97 \pm 0.08	1.1 \pm 0.5
1000 (M)	3.5 \pm 0.8	4.1 \pm 0.9	1.2 \pm 0.03	1.1 \pm 0.4

^a n.e = not estimated

Table 5

The analyzed concentrations of TNT and TNT metabolites in the bile. The results are mean concentrations (mg/L \pm SD) in bile from 4 fish. The medians are shown in brackets. M= water from Lake Maridalsvannet

Exposure ($\mu\text{g/L}$)	TNT (mg/L)	2-ADNT (mg/L)	4-ADNT (mg/L)
1	< 2	< 2	< 2
10	< 2	1.1 \pm 0.02 (0.7)	5.5 \pm 3.8 (4.5)
100	< 2	12 \pm 12 (6.8)	40 \pm 26 (28)
1000^a	< 2	29 \pm 20 (36)	50 \pm 33 (67)
1000 (M)	< 2	60 \pm 42 (55)	56 \pm 37 (53)

^a Results from 3 fish

Table 6

Effects on blood physiology parameters of the fish exposed to TNT. The results are presented as mean concentration \pm SD. The medians are shown in the brackets. Each group was compared to the control group (Ctr) by one-way ANOVA. * represent a significance level of $P < 0.05$, ***represent a significance level of $P < 0.001$.

Exposure ($\mu\text{g/L}$)	HCO₃ (mmol/L)	Na (mmol/L)	K (mmol/L)	Cl (mmol/L)	n
0 (Ctr)	3.4 \pm 0.8 (3.2)	147 \pm 2.9 (147)	2.9 \pm 1.2 (2.3)	134 \pm 1.1 (134)	5
0 (Solvent)	3.7 \pm 0.7 (3.5)	145 \pm 2.6 (146)	4.8 \pm 1.2 (4.9)	133 \pm 3.0 (134)	6
1	3.5 \pm 0.9 (3.2)	145 \pm 3.4 (146)	5.2 \pm 1.1 (4.9)	136 \pm 1.1 (136)	6
10	3.8 \pm 1.0 (3.9)	144 \pm 2.0 (145)	5.3 \pm 1.0 (5.3)	135 \pm 3.1 (135)	6
100	2.9 \pm 0.4 (3.1)	141 \pm 2.8 (141)	4.9 \pm 0.7 (5.0)	134 \pm 1.4 (134)	6
1000	9.5 \pm 3.8 (8.3)***	137 \pm 2.1 (138)	4.3 \pm 0.4 (4.3)	115 \pm 2.1 (115)***	6
1000 (M)	3.0 \pm 1.0 (2.5)	141 \pm 4.3 (139)	5.9 \pm 1.4 (5.5)	137 \pm 2.8 (138)	5

Table 7

Effects on blood physiology parameters of the fish exposed to TNT. The results are presented as mean concentrations \pm SD. The medians are shown in the brackets. Each group was compared with the control group (Ctr) by one-way ANOVA. * represent a significance level of $P < 0.05$, ***represent a significance level of $P < 0.001$.

Exposure ($\mu\text{g/L}$)	Glucose (mmol/L)	Urea (mg/dL)	Hct (% PCV)	Hb (g/dL)	n
0 (Ctr)	3.7 \pm 0.8 (3.9)	< 1	42 \pm 3.4 (42)	14.1 \pm 1.1 (14.3)	5
0 (Solvent)	4.7 \pm 1.3 (4.8)	< 1	36 \pm 3.3 (37)	12.1 \pm 1.1 (12.6)	6
1	4.4 \pm 1.1 (4.6)	< 1	37 \pm 3.0 (37)	12.4 \pm 1.0 (12.4)	6
10	3.5 \pm 0.8 (3.5)	< 1	35 \pm 5.6 (35)*	11.8 \pm 1.9 (11.9)	6
100	3.7 \pm 0.3 (3.8)	< 1	37 \pm 2.7 (37)	12.4 \pm 0.9 (12.6)	6
1000	14 \pm 2.7 (13)***	1.5 \pm 0.5 (1.3)***	16 \pm 3.3 (15)***	5.4 \pm 1.1 (5.1)***	6
1000 (M)	11 \pm 3.0 (12)***	2.6 \pm 0.5 (2.8)***	20 \pm 2.6 (19)***	6.8 \pm 0.9 (6.5)***	5




Fig 1. Measured radioactivity in the water shown as $\mu\text{g/L } ^{14}\text{C-TNT}$ equivalents. After 48 hours the fish were transferred to clean water showing an increase in radioactivity, probably due to excretion from the fish.




Fig 2. Uptake and excretion of $^{14}\text{C-TNT}$ as a function of time in different organs. After 48 h the fish were transferred to clean water showing depuration of the $^{14}\text{C-TNT}$. Each data point represent mean (\pm SD) from 6 fish.




Fig 3. Uptake and excretion of ¹⁴C-TNT as a function of time in the intestine and gall bladder. After 48h the fish were transferred to clean water showing depuration of the ¹⁴C-TNT. Each data point represent mean (\pm SD) from 6 fish.

Supplementary materials

Uptake and effects of 2, 4, 6 - trinitrotoluene (TNT) in juvenile Atlantic salmon
(*Salmo salar*)

Espen Mariussen,^{1,2} Siv Marie Stornes³, Kari Oline Bøifot¹, Bjørn Olav Rosseland³, Brit Salbu,³ and Lene Sørli Heier^{3,4}

¹Norwegian Defence Research Establishment (FFI), P.O. Box 25, N-2027 Kjeller, Norway

²Norwegian Institute for Air Research (NILU), P.O. Box 100 NO-2027 Kjeller, Norway

³Norwegian University of Life Sciences (NMBU), Faculty of Environmental Sciences and Natural Resource Management, P.O. Box 5003, N-1432 Ås, Norway

⁴Norwegian Public Roads Administration, Region Øst, P.O. Box 1010 Nordre Ål, 2605 Lillehammer, Norway

Correspondence: Espen Mariussen

Norwegian Institute for Air Research (NILU),

P. O. Box 100, N-2027 Kjeller, Norway

Phone: + 47 63 89 80 00

Fax: + 47 63 89 80 50 E-mail: Espen.Mariussen@nilu.no




Fig 1S. Estimated elimination half-time of ^{14}C -TNT in different organs collected from the fish from the time the depuration started. Each data point represent the mean from six fish \pm SD. Elimination half-time is estimated from non-linear regression one-phase decay. The results after 48 h represents retained radioactivity in the fish after transfer to clean water



Fig 2S. Juvenile Salmon exposed to 1 mg/L TNT with hemorrhages in the dorsal muscle tissue near the spine.